

DURAMUNE® MAX 5-CvK/4L THE PUPPYSHOT® BOOSTER

Boehringer Ingelheim

Canine Distemper-Adenovirus Type 2-Coronavirus-Parainfluenza-Parvovirus Vaccine

Modified Live and Killed Virus

Leptospira Canicola-Grippytyphosa-Icterohaemorrhagiae-Pomona Bacterial Extract

For subcutaneous vaccination of healthy dogs 6 weeks of age or older as an aid in the prevention of disease caused by canine distemper, infectious canine hepatitis, canine adenovirus type 2, canine coronavirus, canine parainfluenza, canine parvovirus, *Leptospira canicola*, *Leptospira icterohaemorrhagiae*, *Leptospira grippytyphosa* and *Leptospira pomona*.

DIRECTIONS FOR USE

General Directions: Aseptically rehydrate DURAMUNE Max 5 with Canine Coronavirus vaccine-Leptospira bacterial extract supplied. Administer one 1 mL dose subcutaneously.

Primary Vaccination: A recommended vaccination schedule should start at or about 6 weeks of age. The presence of maternal antibody is known to interfere with the development of active immunity. Puppies should be revaccinated every 2 to 3 weeks until 12 weeks of age.

All dogs over 12 weeks of age should initially receive one dose of Duramune Max 5-CvK/4L and a second dose 2 to 3 weeks later.

Annual Vaccination: Annual revaccination with one dose is recommended.